

# An Erdős-Ko-Rado Theorem For Signed Sets

B. BOLLOBÁS<sup>†</sup>

Department of Mathematics  
 University of Memphis, Memphis, TN 38152, U.S.A.  
 and

Department of Pure Mathematics and Mathematical Statistics  
 University of Cambridge, England

I. LEADER

Department of Pure Mathematics and Mathematical Statistics  
 University of Cambridge, England

**Abstract**—A *signed  $r$ -set* on  $[n] = \{1, \dots, n\}$  is a pair  $(A, f)$ , where  $A \subset [n]$  is an  $r$ -set and  $f$  is a function from  $A$  to  $\{-1, 1\}$ . A family  $\mathcal{A}$  of signed  $r$ -sets is *intersecting* if for any  $(A, f), (B, g) \in \mathcal{A}$  there exists  $x \in A \cap B$  such that  $f(x) = g(x)$ . In this note, we prove that if  $\mathcal{A}$  is an intersecting family of signed  $r$ -sets on  $[n]$ , then  $|\mathcal{A}| \leq 2^{r-1} \binom{n-1}{r-1}$ .

We also present an application of this result to a diameter problem in the grid.

**Keywords**—Extremal problems, Intersecting families, Discrete isoperimetric inequalities.

## 1. AN ERDŐS-KO-RADO THEOREM FOR SIGNED SETS

A *signed  $r$ -set* on  $[n] = \{1, \dots, n\}$  is a pair  $(A, f)$ , where  $A \subset [n]$  is an  $r$ -set and  $f$  is a function from  $A$  to  $\{-1, 1\}$ . A family  $\mathcal{A}$  of signed  $r$ -sets is *intersecting* if for any  $(A, f), (B, g) \in \mathcal{A}$  there exists  $x \in A \cap B$  such that  $f(x) = g(x)$ .

Our aim is to prove that if  $\mathcal{A}$  is an intersecting family of signed  $r$ -sets on  $[n]$  then  $|\mathcal{A}| \leq 2^{r-1} \binom{n-1}{r-1}$ . Note that this inequality is best possible: writing  $S_{n,r}$  for the collection of all signed  $r$ -sets on  $[n]$ , for any  $x_0 \in [n]$  and  $\epsilon_0 \in \{-1, 1\}$ , the family  $\{(A, f) \in S_{n,r} : x_0 \in A, f(x_0) = \epsilon_0\}$  is an intersecting family of signed  $r$ -sets of size  $2^{r-1} \binom{n-1}{r-1}$ .

Note also that if  $\mathcal{A}$  is an intersecting family of signed  $r$ -sets on  $[n]$  then, for each  $r$ -set  $A$ , there can be at most  $2^{r-1}$  functions  $f : A \rightarrow \{-1, 1\}$  such that  $(A, f) \in \mathcal{A}$ . Furthermore, the system of underlying  $r$ -sets of  $\mathcal{A}$ , namely  $\{A : (A, f) \in \mathcal{A} \text{ for some } f\}$ , is an intersecting family. Thus  $|\mathcal{A}|$  is at most  $2^{r-1}$  times the maximal size of an intersecting family of  $r$ -sets on  $[n]$ . In particular, if  $r \leq n/2$  then by the Erdős-Ko-Rado Theorem [1], the maximal size of an intersecting family of  $r$ -sets is  $\binom{n-1}{r-1}$ , and so  $|\mathcal{A}| \leq 2^{r-1} \binom{n-1}{r-1}$ . Thus, the interest of our result is in the case  $r > n/2$ .

Our method of proof is an averaging method, similar to that used by Katona [2] in his beautiful proof of the Erdős-Ko-Rado Theorem itself. The proof of the inequality itself is rather short, but we have to work a little to prove that the extremal systems are essentially unique.

**THEOREM 1.** *Let  $1 \leq r \leq n$ , and let  $\mathcal{A}$  be an intersecting family of signed  $r$ -sets on  $[n]$ . Then  $|\mathcal{A}| \leq 2^{r-1} \binom{n-1}{r-1}$ . If  $r < n$ , then equality holds if and only if  $\mathcal{A} = \{(A, f) \in S_{n,r} : x_0 \in A, f(x_0) = \epsilon_0\}$  for some  $x_0 \in [n]$  and  $\epsilon_0 \in \{-1, 1\}$ .*

<sup>†</sup>Research partially supported by NSF Grant DMS-8806097.

PROOF. Call a cyclic ordering of  $[n] \times \{-1, 1\}$  *good* if  $(x, 1)$  is opposite  $(x, -1)$  for all  $x \in [n]$ . More formally, a cyclic ordering is a bijection  $\sigma : [n] \times \{-1, 1\} \rightarrow \mathbb{Z}_{2n}$ , and  $\sigma$  is said to be good if  $\sigma(x, 1) = \sigma(x, -1) + n$  for all  $x \in [n]$ . We say that a signed  $r$ -set  $(A, f)$  is *compatible* with a cyclic ordering if there is an interval  $(x_1, \epsilon_1), \dots, (x_r, \epsilon_r)$  in the ordering such that for each  $i$  we have  $x_i \in A$  and  $f(x_i) = \epsilon_i$ . In other words,  $(A, f)$  is compatible with  $\sigma$  if  $A$  can be written as  $\{x_1, \dots, x_r\}$  such that, for some  $z \in \mathbb{Z}_{2n}$ , we have  $\sigma(x_i, f(x_i)) = z + i$  for all  $i$ .

Fix a good cyclic ordering  $\sigma$ . How many members of  $\mathcal{A}$  are compatible with  $\sigma$ ? The corresponding intervals must intersect pairwise. However, since  $r \leq (1/2) \cdot 2n$ , it is easy to see that at most  $r$  intervals of length  $r$  may intersect pairwise. Thus, at most  $r$  members of  $\mathcal{A}$  are compatible with  $\sigma$ .

Now, there are  $n! 2^n$  good cyclic orderings, and any fixed signed  $r$ -set  $(A, f)$  is compatible with exactly  $2n \cdot r! (n - r)! 2^{n-r}$  of them. It follows that

$$2n \cdot r! (n - r)! 2^{n-r} |\mathcal{A}| \leq r \cdot 2^n n!,$$

and so  $|\mathcal{A}| \leq \binom{n-1}{r-1} 2^{r-1}$ , proving our inequality.

Let us turn to the case of equality: suppose that  $r < n$  and  $|\mathcal{A}| = \binom{n-1}{r-1} 2^{r-1}$ . Then each good cyclic ordering  $\sigma$  is compatible with exactly  $r$  members of  $\mathcal{A}$ . The corresponding intervals intersect pairwise, and so, since  $r < (1/2) \cdot 2n$ , they must consist of all intervals containing a fixed element of  $[n] \times \{-1, 1\}$ ; we denote this fixed element by  $(x^{(\sigma)}, \epsilon^{(\sigma)})$ . Note that, in particular, whenever there are two consecutive intervals  $(x_1, \epsilon_1), \dots, (x_r, \epsilon_r)$  and  $(x_2, \epsilon_2), \dots, (x_{r+1}, \epsilon_{r+1})$  in the ordering  $\sigma$ , such that the first belongs to  $\mathcal{A}$  but the second does not, then the fixed element  $(x^{(\sigma)}, \epsilon^{(\sigma)})$  must be precisely  $(x_1, \epsilon_1)$ .

Let  $\tau$  be the good cyclic ordering given by

$$\tau(x, \epsilon) = \begin{cases} x, & \text{if } \epsilon = 1, \\ x + n, & \text{if } \epsilon = -1, \end{cases}$$

and assume without loss of generality that  $(x^{(\tau)}, \epsilon^{(\tau)}) = (n, 1)$ . Given  $1 \leq i < j \leq n - 1$ , let  $\tau_{ij}$  be the good cyclic ordering obtained from  $\tau$  by swapping  $i$  and  $j$ :

$$\tau_{ij}(x, \epsilon) = \begin{cases} \tau(j, \epsilon), & \text{if } x = i, \\ \tau(i, \epsilon), & \text{if } x = j, \\ \tau(x, \epsilon), & \text{otherwise.} \end{cases}$$

Then, using the remark at the end of the previous paragraph, it is routine to check that we must have  $(x^{(\tau_{ij})}, \epsilon^{(\tau_{ij})}) = (x^{(\tau)}, \epsilon^{(\tau)}) = (n, 1)$ .

Since  $i$  and  $j$  were arbitrary, it follows that if  $\pi$  is any permutation of  $[n - 1]$ , and we define the cyclic ordering  $\tau_\pi$  by

$$\tau_\pi(x, \epsilon) = \begin{cases} \tau(\pi(x), \epsilon), & \text{if } x < n, \\ \tau(x, \epsilon), & \text{if } x = n, \end{cases}$$

then also  $(x^{(\tau_\pi)}, \epsilon^{(\tau_\pi)}) = (n, 1)$ . However, it is clear that if  $(A, f)$  is any signed  $r$ -set on  $[n]$  with  $n \in A$  and  $f(n) = 1$ , then  $(A, f)$  corresponds to an interval in the ordering  $\tau_\pi$  for some  $\pi$ . Indeed, if  $A = \{n, x_1, \dots, x_s, y_1, \dots, y_{r-s-1}\}$ , with  $f(n) = 1$ ,  $f(x_i) = 1$  for all  $i$  and  $f(y_j) = -1$  for all  $j$ , then we may take any permutation  $\pi$  satisfying  $\pi(x_1) = n - 1, \pi(x_2) = n - 2, \dots, \pi(x_s) = n - s, \pi(y_1) = 1, \pi(y_2) = 2, \dots, \pi(y_{r-s-1}) = r - s - 1$ . It follows that  $\mathcal{A}$  contains  $\{(A, f) \in S_{n,r} : n \in A, f(n) = 1\}$ , as required.  $\blacksquare$

It is fairly simple to extend Theorem 1 to signed sets which may take more than two values. An  $l$ -signed  $r$ -set is a pair  $(A, f)$ , where  $A \subset [n]$  is an  $r$ -set and  $f$  is a function from  $A$  to  $[l]$ . We write  $S_{n,r,l}$  for the collection of all  $l$ -signed  $r$ -sets on  $[n]$ . A family  $\mathcal{A}$  of  $l$ -signed  $r$ -sets is *intersecting* if for any  $(A, f), (B, g) \in \mathcal{A}$  there exists  $x \in A \cap B$  such that  $f(x) = g(x)$ .

**THEOREM 2.** Let  $1 \leq r \leq n$ , and let  $\mathcal{A}$  be an intersecting family of  $l$ -signed  $r$ -sets on  $[n]$ , where  $l \geq 2$ . Then  $|\mathcal{A}| \leq \binom{n-1}{r-1} l^{r-1}$ . Unless  $l = 2$  and  $r = n$ , equality holds if and only if  $\mathcal{A} = \{(A, f) \in S_{n,r,l} : x_0 \in A, f(x_0) = i_0\}$  for some  $x_0 \in [n]$  and  $i_0 \in [l]$ .

**PROOF.** Call a cyclic ordering  $\sigma$  of  $[n] \times [l]$  (a bijection from  $[n] \times [l]$  to  $\mathbb{Z}_{nl}$ ) *good* if for every  $x \in [n]$  and  $i \in [l]$  there is a  $j \in [l]$  with  $\sigma(x, i) + n = \sigma(x, j)$ . We now proceed just as in the proof of Theorem 1.

Note that, in the case of equality, for  $l \geq 3$  we do not need the restriction  $r < n$ , since if  $l \geq 3$  then any  $r \leq n$  satisfies  $r < (1/2)kn$ . ■

We may reformulate Theorem 2 as follows.

**THEOREM 2'.** Let  $l \geq 2$ , and let  $X = X_1 \cup \dots \cup X_n$ , where the  $X_i$  are disjoint sets each of size  $l$ . Let  $1 \leq r \leq n$ , and let  $\mathcal{A}$  be an intersecting family of  $r$ -sets on  $X$  such that every  $A \in \mathcal{A}$  satisfies  $|A \cap X_i| \leq 1$  for all  $i$ . Then  $|\mathcal{A}| \leq \binom{n-1}{r-1} l^{r-1}$ . Moreover, except in the case  $l = 2, r = n$ , equality holds if and only if all members of  $\mathcal{A}$  contain a fixed element  $x_0$  of  $X$ . ■

## 2. AN APPLICATION

Let  $k$  be a positive integer, and write  $[-k, k]$  for  $\{-k, -k+1, \dots, k\}$ . The *grid* on  $[-k, k]^n$  is the graph on  $[-k, k]^n$  in which  $(x_1, \dots, x_n)$  is joined to  $(y_1, \dots, y_n)$ , if for some  $i$  we have  $|x_i - y_i| = 1$  and  $x_j = y_j$  for all  $j \neq i$ . Given a positive integer  $d$ , how large can a subset of  $[-k, k]^n$  of diameter  $d$  be?

For any  $c \in \mathbb{R}^n$  and  $r \in \mathbb{R}_+$  let us set

$$B(c, r) = \left\{ x \in [-k, k]^n : \sum |x_i - c_i| \leq r \right\}.$$

Thus, certainly  $B(c, r)$  has diameter at most  $2r$ . Kleitman and Fellows [3] conjectured that if  $d$  is even, then a subset of  $[-k, k]^n$  of diameter  $d$  has size at most  $|B(0, d/2)|$ , while if  $d$  is odd then the maximum size is  $|B((1/2)e_1, d/2)|$  (where  $e_1, \dots, e_n$  denotes the standard basis of  $\mathbb{R}^n$ ). This was proved in [4].

Our aim here is to give a different proof of this result when  $d \geq n(k+1)$ , in other words for values of  $d$  larger than ‘half-way’. Not surprisingly, the case  $d$  odd is harder than the case  $d$  even, and it is there that we shall apply Theorem 1.

We need a small amount of notation. Let  $G$  be a subset of  $[-k, k]^n$ , and for convenience write  $S_r$  for  $S_{n,r}$ . For  $(A, f) \in S_r$ , we define the  $(A, f)$ -*quadrant* of  $G$  to be

$$G_{(A,f)} = \left\{ x \in [k]^A : \sum_{i \in A} x_i f(i) e_i \in G \right\}.$$

Thus,  $|G_{(A,f)}|$  is the number of points  $x \in G$  such that  $x_i = 0$  for  $i \notin A$ , and for  $i \in A$  the sign of  $x_i$  is  $f(i)$ . In particular,

$$|G| \leq 1 + \sum_{r=1}^n \sum_{(A,f) \in S_r} |G_{(A,f)}|,$$

with equality if  $0 \in G$ .

For  $1 \leq r \leq n$ , and any nonnegative integer  $s$ , define

$$F_r(s) = \left| \left\{ x \in [k]^r : \sum x_i = s \right\} \right|.$$

Thus,  $F_r(s) = F_r(r(k+1) - s)$  for all  $s$ . It is easy to check that the function  $F_r$  is increasing for  $s \leq r(k+1)/2$ , for example, because  $[k]^r$  has a symmetric chain decomposition (see [5, Chapter 3]).

We are now ready for our application.

**THEOREM 3.** *Let  $G$  be a subset of  $[-k, k]^n$  of diameter  $d$ , where  $d \geq n(k+1)$ . Then*

$$|G| \leq \begin{cases} \left| B\left(0, \frac{d}{2}\right) \right|, & \text{if } d \text{ is even,} \\ \left| B\left(\frac{1}{2}e_1, \frac{d}{2}\right) \right|, & \text{if } d \text{ is odd.} \end{cases}$$

**PROOF.** Let us start with the case  $d$  even. For convenience, write  $B$  for  $B(0, d/2)$ . To show that  $|G| \leq |B|$ , we shall show that the total number of points in ‘opposite quadrants’ is at most as large for  $G$  as it is for  $B$ ; in other words, we shall show that for every  $1 \leq r \leq n$  and every  $(A, f) \in S_r$ , we have

$$|G_{(A,f)}| + |G_{(A,-f)}| \leq |B_{(A,f)}| + |B_{(A,-f)}|.$$

Fix then a signed  $r$ -set  $(A, f)$ . If either of  $G_{(A,f)}$ ,  $G_{(A,-f)}$  is empty then certainly  $|G_{(A,f)}| + |G_{(A,-f)}| \leq |B_{(A,f)}| + |B_{(A,-f)}|$ , since the fact that  $d \geq r(k+1)$  implies that  $|B_{(A,f)}| + |B_{(A,-f)}| \geq k^n$ . So we may assume that  $G_{(A,f)}, G_{(A,-f)} \neq \emptyset$ .

Choose  $x \in G_{(A,f)}$  and  $y \in G_{(A,-f)}$  with  $\sum x_i$  and  $\sum y_i$  maximal: say  $\sum x_i = t$  and  $\sum y_i = u$ . Since  $G$  has diameter  $d$ , we must have  $t + u \leq d$ . Now, by the choice of  $t$  and  $u$ , we have

$$|G_{(A,f)}| + |G_{(A,-f)}| \leq \sum_{s=r}^t F_r(s) + \sum_{s=r}^u F_r(s).$$

Since  $t + u \leq d$  and  $d \geq r(k+1)$ , we have

$$\sum_{s=r}^t F_r(s) + \sum_{s=r}^u F_r(s) \leq 2 \sum_{s=r}^{d/2} F_r(s) = |B_{(A,f)}| + |B_{(A,-f)}|.$$

Hence  $|G_{(A,f)}| + |G_{(A,-f)}| \leq |B_{(A,f)}| + |B_{(A,-f)}|$ , as required.

We now turn to the case  $d$  odd. Write  $B$  for  $B((1/2)e_1, d/2)$ . For any  $1 \leq r \leq n$  and any  $(A, f) \in S_r$ , it follows just as above that

$$|G_{(A,f)}| + |G_{(A,-f)}| \leq \sum_{s=r}^{(d+1)/2} F_r(s) + \sum_{s=r}^{(d-1)/2} F_r(s).$$

Also, it is certainly true that if for all  $x \in G_{(A,f)}$  and all  $x \in G_{(A,-f)}$ , we have  $\sum x_i \leq (d-1)/2$  then

$$|G_{(A,f)}| + |G_{(A,-f)}| \leq 2 \sum_{s=r}^{(d-1)/2} F_r(s).$$

Now,  $B_{(A,f)} = \{x \in [k]^A : \sum x_i \leq (d+1)/2\}$  if  $1 \in A$  and  $f(1) = 1$ , and  $\{x \in [k]^A : \sum x_i \leq (d-1)/2\}$  otherwise. Thus, to complete the proof that  $|G| \leq |B|$ , it will suffice to show that, for each  $1 \leq r \leq n$ , the number of signed  $r$ -sets  $(A, f)$  such that  $\sum x_i \geq (d+1)/2$  for some  $x \in G_{(A,f)}$  is at most  $2^{r-1} \binom{n-1}{r-1}$ .

Let then

$$\mathcal{A} = \left\{ (A, f) \in S_r : \sum x_i \geq \frac{(d+1)}{2} \text{ for some } x \in G_{(A,f)} \right\}.$$

We claim that  $\mathcal{A}$  is an intersecting family. Indeed, suppose that, to the contrary, there are signed  $r$ -sets  $(A, f), (C, g) \in \mathcal{A}$  such that for all  $i \in A \cap C$  we have  $f(i) \neq g(i)$ . Then, for any  $x \in G_{(A,f)}$  and  $y \in G_{(C,g)}$ , the distance between  $\sum x_i f(i) e_i$  and  $\sum y_i g(i) e_i$  is  $\sum x_i + \sum y_i$ . However, this implies that  $\mathcal{A}$  has diameter at least  $d+1$ .

Thus,  $\mathcal{A}$  is an intersecting family of signed  $r$ -sets on  $[n]$ , and so by Theorem 1 we have  $|\mathcal{A}| \leq 2^{r-1} \binom{n-1}{r-1}$ , as required.  $\blacksquare$

We remark that it is not hard to see that if equality holds in Theorem 3, and  $d$  is even, then in fact  $G = B(0, d/2)$ . Also, using the case of equality in Theorem 1, one may check that if equality holds in Theorem 3 and  $d$  is odd then  $G = B(\{-1, 1\}(1/2)e_i, (d/2))$  for some  $i$  and choice of sign.

As we mentioned above, it was proved in [4] that the conclusion of Theorem 3 remains valid without the restriction that  $d \geq n(k+1)$ . The same paper also contains several related results.

## REFERENCES

1. P. Erdős, C. Ko and R. Rado, Intersection theorems for systems of finite sets, *Quart. J. Math. Oxford* **12** (2), 313–320 (1961).
2. G.O.H. Katona, A simple proof of the Erdős-Ko-Rado theorem, *J. Combinatorial Theory (B)* **13**, 183–184 (1972).
3. D.J. Kleitman and M. Fellows, Radius and diameter in Manhattan lattices, *Disc. Math.* **73**, 119–125 (1988).
4. B. Bollobás and I. Leader, Maximal sets of given diameter in the grid and the torus, *Disc. Math.* **122**, 15–35 (1993).
5. I. Anderson, *Combinatorics of Finite Sets*, Oxford University Press, pp. xv–250, (1987).